arXiv:1406.1654v2 [math.PR] 12 Nov 2014

On the Moment Determinacy of Products of Non-identically Distributed Random Variables

Gwo Dong Lin[1] • **Jordan Stoyanov**[2]

[1] Institute of Statistical Science, Academia Sinica, Taipei 11529, Taiwan (ROC)
e-mail: gdlin@stat.sinica.edu.tw

[2] School of Mathematics & Statistics, Newcastle University, Newcastle upon Tyne NE1 7RU, UK
e-mail: stoyanovj@gmail.com

Abstract: We show first that there are intrinsic relationships among different conditions, old and recent, which lead to some general statements in both the Stieltjes and the Hamburger moment problems. Then we describe checkable conditions and prove new results about the moment (in)determinacy for products of independent and non-identically distributed random variables. We treat all three cases when the random variables are nonnegative (Stieltjes case), when they take values in the whole real line (Hamburger case), and the mixed case. As an illustration we characterize the moment determinacy of products of random variables whose distributions are generalized gamma or double generalized gamma all with distinct shape parameters. Among other corollaries, the product of two independent random variables, one exponential and one inverse Gaussian, is moment determinate, while the product is moment indeterminate for the cases: one exponential and one normal, one chi-square and one normal, and one inverse Gaussian and one normal.

Mathematics Subject Classification (2010) 60E05, 44A60

Keywords Product of random variables, Stieltjes moment problem, Hamburger moment problem, Cramér's condition, Carleman's condition, Krein's condition, Hardy's condition.

1. Introduction

There is a long standing interest in studying products of random variables; see, e.g., [13], [16], [22], [6], [21], [10], [26] and the references therein. The reasons are two-fold. On one hand, to deal with products leads to non-trivial, difficult and challenging theoretical problems requiring to use diverse ideas and techniques. Let us mention just a few sources: [10], [1], [3]. On the other hand, products of random variables are naturally involved in stochastic modelling of complex random phenomena in areas such as statistical physics, quantum theory, communication theory and financial modelling; see, e.g., [4], [9], [10], [7], [12], [8], [20], [5].

In general, it is difficult to find explicit closed-form expressions for the densities or the distributions of products of random variables with different distributions. It is, however, possible to study successfully the moment problem for products of independent random variables; see, e.g., [15], [25]. Answers about the moment (in)determinacy can be found if requiring only information about the asymptotic of the moments or about the tails of the densities or of their distributions.

All random variables considered in this paper are defined on an underlying probability space $(\Omega, \mathcal{F}, \mathbf{P})$ and we denote by $\mathbf{E}[X]$ the expected value of the random variable X. A basic assumption is that the random variables we deal with have finite moments of all positive orders, i.e. $\mathbf{E}[|X|^k] < \infty,\ k = 1, 2, \ldots$. We write $X \sim F$ to mean that X is a random variable whose distribution function is F and denote its kth order moment by $m_k = \mathbf{E}[X^k]$. We say that X or F is moment determinate (M-det) if F is the only distribution having the moment sequence $\{m_k\}_{k=1}^{\infty}$; otherwise, we say that X or F is moment indeterminate (M-indet). We use traditional notions, notations and terms such as Cramér's condition, Carleman's condition, Krein's condition, and Hardy's condition. For reader's convenience we give their definitions in the text.

We use $\Gamma(\cdot)$ for the Euler-gamma function, $\mathbb{R} = (-\infty, \infty)$ for the set of all real numbers, $\mathbb{R}_+ = [0, \infty)$ for the nonnegative numbers, the symbol $\mathcal{O}(\cdot)$ with its usual meaning in asymptotic analysis and the abbreviation i.i.d. for independent and identically distributed (random variables).

In Section 2 we describe useful intrinsic relationships among different old and recent conditions involved in the Stieltjes and/or the Hamburger moment problems. Then we prove some new results under conditions which are relatively easy to check. In Section 3 we deal with the moment determinacy of products of independent nonnegative random variables with different distributions, while in Section 4 we consider products of random variables with values in $\mathbb{R}$. Finally, in Section 5, we treat the mixed case: products of both types of random variables, nonnegative ones and real ones, the latter with values in $\mathbb{R}$.

The results presented in this paper extend some previous results for products of i.i.d. random variables. Here we need a more refined analysis of the densities of products than in the i.i.d. case. As an illustration we characterize the moment (in)determinacy of products of random variables whose distributions are generalized gamma or double generalized gamma all with distinct shape parameters. We have derived several corollaries involving popular distributions widely used in theoretical studies and applications. Let us list a few: (i) the product of two independent random variables, one exponential and one inverse Gaussian, is M-det; (ii) the product of independent exponential and normal random variables is M-indet; (iii) the product of independent chi-square and normal random variables is M-indet; and (iv) the product of independent inverse Gaussian and normal random variables is M-indet.

2. Some General Results

In this section we present two lemmas, each containing workable conditions which, more or less, are available in the literature. Some of these conditions are old, while others are recent. We describe intrinsic relationships among these conditions and use them to prove new results; see Theorems 1–4.

Our findings in this section can be considered as a useful complement to the classical criteria of Cramér, Carleman, Krein, Hardy and their converses, so that all these taken together make more clear, and possibly complete, the picture of what is in our hands when discussing the determinacy of distributions in terms of their moments.

2.1. Stieltjes Case

Lemma 1. *Let $0 \le X \sim F$. Then the following four statements are equivalent:*
(i) $m_k = \mathcal{O}(k^{2k})$ as $k \to \infty$.
(ii) $\limsup_{k\to\infty} \frac{1}{k}\, m_k^{1/(2k)} < \infty$.
(iii) $m_k \le c_0^k\,(2k)!,\ k = 1, 2, \ldots$, for some constant $c_0 > 0$.
(iv) X satisfies Hardy's condition, namely, $\mathbf{E}[e^{c\sqrt{X}}] < \infty$ for some constant $c > 0$.

The equivalence of conditions (i) and (ii), a known fact for decades, can be easily checked. Conditions (iii) and (iv) appeared recently and their equivalence to condition (ii) were shown in [24].

Theorem 1. *Let $0 \le X \sim F$ with moments growing as follows: $m_k = \mathcal{O}(k^{ak})$ as $k \to \infty$ for some constant $a \in (0, 2]$. Then the following two statements hold:*
(i) X satisfies Hardy's condition and hence X is M-det.
(ii) The boundary value $a = 2$ is the best possible for X to be M-det. In fact, there is an M-indet random variable $\tilde{X} \ge 0$ such that $\mathbf{E}[\tilde{X}^k] = \mathcal{O}(k^{ak})$ as $k \to \infty$ for all $a > 2$.

Proof. Part (i) follows easily from Lemma 1. To prove part (ii), we consider the following absolutely continuous distribution $\tilde{F}$ whose density is

$$\tilde{f}(x) = \tilde{c}\, e^{-\sqrt{x}/(1+|\ln x|^{\delta})}, \quad x > 0. \tag{1}$$

Here $\delta > 1$ and $\tilde{c}$ is the norming constant. Then it is easy to evaluate the Krein quantity for $\tilde{X} \sim \tilde{F}$. Recall that $\tilde{X}$ is nonnegative, so in this Stieltjes case we obtain

$$\mathrm{K}[\tilde{f}] := \int_0^{\infty} \frac{-\ln \tilde{f}(x^2)}{1 + x^2} dx < \infty.$$

Hence $\tilde{X}$ is M-indet (see, e.g., [14], Theorem 3).

The next step is to check that $\mathbf{E}[\tilde{X}^k] = \mathcal{O}(k^{ak})$ as $k \to \infty$ for all $a > 2$. To see this, we fix $a > 2$, take $b \in (2, a)$ and easily find a number $x_0 \ge 1$ such that

$\sqrt{x} > x^{1/b}(1 + |\ln x|^{\delta})$ for all $x \geq x_0$. We now have that

$$\int_0^{x_0} x^k \tilde{f}(x)dx \leq \frac{\tilde{c}}{k+1}\, x_0^{k+1} = \mathcal{O}(k^{ak}) \;\text{ as }\; k \to \infty$$

and that

$$\begin{aligned}\int_{x_0}^{\infty} x^k \tilde{f}(x)dx &\leq \tilde{c}\int_{x_0}^{\infty} x^k e^{-x^{1/b}}dx \leq \tilde{c}\int_0^{\infty} x^k e^{-x^{1/b}}dx \\ &= b\,\tilde{c}\,\Gamma((k+1)b) = \mathcal{O}(k^{ak}) \;\text{ as }\; k \to \infty.\end{aligned}$$

For the last relation we have used the approximation of the gamma function:

$$\Gamma(x) \approx \sqrt{2\pi}\, x^{x-1/2}\, e^{-x} \quad \text{as} \;\; x \to \infty$$

(see, e.g., [27], p. 253). Thus we have shown that indeed $\mathbf{E}[\tilde{X}^k] = \mathcal{O}(k^{ak})$ as $k \to \infty$ for all $a > 2$. Therefore the constant 2 in the formulation of the theorem is indeed the best possible for X to be M-det. □

Remark 1. For $0 \leq X \sim F$, let us compare the following two moment conditions: (a) $m_k = \mathcal{O}(k^{2k})$ as $k \to \infty$, and (b) $m_{k+1}/m_k = \mathcal{O}((k+1)^2)$ as $k \to \infty$. Here (a) is the condition in Theorem 1, while condition (b) was introduced and used in the recent paper [15]. Both conditions are checkable and each of them guarantees the moment determinacy of F. Just to mention that condition (b) implies condition (a) by referring to Theorem 3 in [15], while the converse may not be true in general.

The next result, Theorem 2 below, is the converse to Theorem 1, and deals with the moment indeterminacy of nonnegative random variables. We need first one condition which is used a few times in the sequel.

Condition L: Suppose, in the Stieltjes case, that $f(x),\ x \in \mathbb{R}_+$, is a density function such that for some fixed $x_0 > 0$, f is strictly positive and differentiable for $x > x_0$ and

$$L_f(x) := -\frac{x f'(x)}{f(x)} \nearrow \infty \;\text{ as }\; x_0 < x \to \infty.$$

In the Hamburger case we require the density $f(x),\ x \in \mathbb{R}$, to be symmetric.

This condition plays a significant rôle in moment problems for absolutely continuous probability distributions. It was explicitly introduced and efficiently used for the first time in [14] and later used by several authors naming it as 'Lin's condition'. This condition is involved in some of our results to follow.

Theorem 2. *Let $0 \leq X \sim F$ and its moment sequence $\{m_k,\ k = 1, 2, \ldots\}$ grow 'fast' in the sense that $m_k \geq c\, k^{(2+\varepsilon)k},\ k = 1, 2, \ldots$, for some constants $c > 0$ and $\varepsilon > 0$. Assume further that X has a density function f which satisfies the above Condition L. Then X is M-indet.*

Proof. By the condition on the moments, we see that the Carleman quantity for the moments of F is finite. Indeed, in this Stieltjes case we have:

$$\mathrm{C}[F] = \sum_{k=1}^{\infty} \frac{1}{m_k^{1/(2k)}} \leq \sum_{k=1}^{\infty} \frac{1}{c^{1/(2k)}\, k^{1+\varepsilon/2}} < \infty.$$

However no conclusion can be drawn from this because $\mathrm{C}[F] < \infty$ is only a necessary condition for X to be M-indet. We need other arguments. By applying Condition L and the second part of the proof of Theorem 4 in [15], we finally conclude that indeed X is M-indet. □

Remark 2. To provide one application of Theorem 2, let us consider, for example, the random variable $X = \xi^{2(1+\varepsilon)}$, where $\varepsilon > 0$ and $\xi \sim Exp(1)$, the standard exponential distribution. On one hand, we can use the Krein criterion and show that X is M-indet. On the other hand, X satisfies the moment condition in Theorem 2. And here is the point: instead of applying Krein's condition, we can prove the moment indeterminacy of X by checking that its density f satisfies Condition L. In general, we follow that approach which seems easier, or which is working in the specific case of interest.

2.2. Hamburger Case

We start with Lemma 2 establishing the equivalence of different type of conditions involved to decide whether a distribution on the whole real line $\mathbb{R}$ is M-det. Then we present some new results. Theorem 3 below is a slight modification, in a new light, of a result in [2], p. 92, while Theorem 4 is the converse to Theorem 3.

Lemma 2. *Let X be a random variable taking values in $\mathbb{R}$. Then the following four statements are equivalent:*
(i) $m_{2k} = \mathcal{O}((2k)^{2k})$ as $k \to \infty$.
(ii) $\limsup_{k\to\infty} \frac{1}{2k}\, m_{2k}^{1/(2k)} < \infty$.
(iii) $m_{2k} \leq c_0^k\, (2k)!,\ k = 1, 2, \ldots$, for some constant $c_0 > 0$.
(iv) X satisfies Cramér's condition: its moment generating function exists.

Proof. It is easy to check the equivalence of conditions (i) and (ii). The equivalence of conditions (ii) and (iv) is well-known, but we provide here a simple and instructive proof based on condition (i). Indeed, by Lemma 1 above, condition (i) is equivalent to say that the random variable $Y = X^2$ satisfies Hardy's condition, namely, $\mathbf{E}[e^{c\sqrt{Y}}] = \mathbf{E}[e^{c|X|}] < \infty$ for some constant $c > 0$. The latter, however, means that X itself has a moment generating function. This is exactly the statement (iv). Finally, applying again Lemma 1 to the nonnegative random variable Y, we obtain the equivalence of (ii) and (iii). Therefore, as stated, all four conditions (i) – (iv) are equivalent. □

Remark 3. In [2] the moment condition $m_k = \mathcal{O}(k^k)$ as $k \to \infty$ was used to derive the M-det of X on $\mathbb{R}$. This condition can be replaced by a weaker one, allowing a

'faster' growth of the moments., e.g., $m_k = \mathcal{O}((a(k))^k)$ as $k \to \infty$, where

$$a(k) = k \ln k, \quad \text{or} \quad k (\ln k) (\ln \ln k), \ \text{or} \quad k (\ln k) (\ln \ln k) (\ln \ln \ln k), \ \ldots,$$

and still preserving the M-det property of X. Such a statement can be established by using quasi-analytic functions. We do not give details here.

Theorem 3. *Let $X \sim F$, where F has an unbounded support $supp(F) \subset \mathbb{R}$ and its moments satisfy the condition: $m_{2k} = \mathcal{O}((2k)^{2ak})$ as $k \to \infty$ for some constant $a \in (0, 1]$. Then the following statements hold:*
(i) X satisfies Cramér's condition and hence is M-det.
(ii) The boundary value $a = 1$ is the best possible for X to be M-det. In fact, there is an M-indet random variable $\tilde{X}$ such that $\mathbf{E}[\tilde{X}^{2k}] = \mathcal{O}((2k)^{2ak})$ as $k \to \infty$ for all $a > 1$.

Proof. Part (i) follows immediately from Lemma 2. Part (ii) is essentially given in [2], but we provide here a somewhat general case. Let us consider the distribution $\tilde{F}$ with the following symmetric density (compare this with (1)):

$$\tilde{f}(x) = \bar{c}\, e^{-|x|/(1+|\ln|x||^{\delta})}, \quad x \in \mathbb{R}, \tag{2}$$

where $\delta > 1$, $\tilde{f}(0) = 1$ and $\bar{c}$ is a norming constant. In this, already Hamburger case, the Krein quantity for $\tilde{X} \sim \tilde{F}$, with density $\tilde{f}$ given by (2), can be evaluated and shown to be finite. Namely, we have that

$$\mathrm{K}[\tilde{f}] := \int_{-\infty}^{\infty} \frac{-\ln \tilde{f}(x)}{1 + x^2} dx < \infty.$$

Hence $\tilde{X}$ is M-indet (see, e.g., [14], Theorem 1). However, it is seen that $\mathbf{E}[\tilde{X}^{2k}] = \mathcal{O}((2k)^{2ak})$ as $k \to \infty$ for all $a > 1$. Therefore, the constant 1 in the formulation of the theorem is indeed the best possible for X to be M-det. □

Remark 4. Suppose $X \sim F$ with F having unbounded support, $\mathrm{supp}(F) \subset \mathbb{R}$. We want to compare the following two moment conditions: (a) $m_{2k} = \mathcal{O}((2k)^{2k})$ as $k \to \infty$, and (b) $m_{2(k+1)}/m_{2k} = \mathcal{O}((k+1)^2)$ as $k \to \infty$. Here (a) is the growth of the moments condition in Theorem 3, while condition (b) was introduced and successfully exploited in the recent work [25]. Both conditions are checkable and each of them guarantees the moment determinacy of X and F. Let us mention that condition (b) implies condition (a) by referring to Theorem 2 in [25], while the converse may not in general be true.

Theorem 4. *Suppose that the moments of $X \sim F$ grow 'fast' in the sense that $m_{2k} \geq c(2k)^{2(1+\varepsilon)k}$, $k = 1, 2, \ldots$, for some positive constants c and ε. Assume further that X has a density function f which is symmetric about zero and satisfies the above Condition L. Then X is M-indet.*

Proof. By the condition on the moments, we see that the Carleman quantity for F is finite (we remember that this is a Hamburger case):

$$\mathrm{C}[F] := \sum_{k=1}^{\infty} \frac{1}{m_{2k}^{1/(2k)}} \leq \sum_{k=1}^{\infty} \frac{1}{c^{1/(2k)}\,(2k)^{1+\varepsilon}} < \infty.$$

Since no conclusion can be drawn from this finding, we need different arguments. We use Condition L and the second part of the proof of Theorem 3 in [25] thus arriving at the desired conclusion that indeed X is M-indet. □

Remark 5. For example, instead of applying Krein's condition, we can use Theorem 4 to prove the moment indeterminacy of $X \sim F$ whose density is the symmetrization of that of $\xi^{1+\varepsilon}$, where $\varepsilon > 0$ and $\xi \sim Exp(1)$.

3. Products of Nonnegative Random Variables

We start with two results describing relatively simple conditions on the random variables $\xi_1, \ldots, \xi_n$ in order to guarantee that their product is M-det.

Theorem 5. *Suppose that the moments $m_{i,k} = \mathbf{E}[\xi_i^k],\ i = 1, \ldots, n,$ of the independent random variables $\xi_1, \ldots, \xi_n$ satisfy the conditions:*

$$m_{i,k} = \mathcal{O}(k^{a_i k}) \quad as \quad k \to \infty, \quad for \quad i = 1, \ldots, n,$$

where $a_1, \ldots, a_n$ are positive constants. If $a_1, \ldots, a_n$ are such that $a_1 + \cdots + a_n \leq 2$, then the product $Z_n = \xi_1 \cdots \xi_n$ is M-det.

Proof. With $m_k = \mathbf{E}[Z_n^k]$ we have, by the independence of ξ_i, that

$$m_k = m_{1,k} \cdots m_{n,k} = \mathcal{O}(k^{a_1 k}) \cdots \mathcal{O}(k^{a_n k}) = \mathcal{O}(k^{ak}) \ \text{ as } \ k \to \infty,$$

where $a = a_1 + \cdots + a_n$. Since, by assumption, $a \leq 2$, we apply Theorem 1 to conclude the M-det property of the product Z_n. □

Theorem 6. *Suppose that the growth rates $r_1, \ldots, r_n$ of the moments of each of the random variables $\xi_1, \ldots, \xi_n$ satisfy*

$$\frac{m_{1,k+1}}{m_{1,k}} = \mathcal{O}((k+1)^{r_1}), \ \ldots, \ \frac{m_{n,k+1}}{m_{n,k}} = \mathcal{O}((k+1)^{r_n}) \quad as \quad k \to \infty,$$

where $m_{i,k} = \mathbf{E}[\xi_i^k],\ i = 1, \ldots, n,\ k = 1, 2, \ldots.$ If the rates $r_1, \ldots, r_n$ are such that $r_1 + \cdots + r_n \leq 2$, then the product $Z_n = \xi_1 \cdots \xi_n$ is M-det.

Proof. Denoting $m_k = \mathbf{E}[Z_n^k]$ and using the independence of ξ_i, we find that

$$\frac{m_{k+1}}{m_k} = \frac{m_{1,k+1}}{m_{1,k}} \cdots \frac{m_{n,k+1}}{m_{n,k}} = \mathcal{O}((k+1)^r) \text{ as } \ k \to \infty,$$

where $r = r_1 + \cdots + r_n$. Since, by assumption, $r \le 2$, we refer to Remark 1 and conclude that the product Z_n is M-det. $\square$

Let us provide now conditions under which the product Z_n becomes M-indet.

Theorem 7. *Consider n independent nonnegative random variables, $\xi_i \sim F_i$, $i = 1, 2, \ldots, n$, where $n \ge 2$. Suppose that each F_i is absolutely continuous with density $f_i > 0$ on $(0, \infty)$ and that the following conditions are satisfied:*
(i) At least one (or just one) of the densities $f_1(x), \ldots, f_n(x)$ is decreasing in $x \ge x_0$, where $x_0 \ge 1$ is a constant.
(ii) For each $i = 1, 2, \ldots, n$, there exists a constant $A_i > 0$ such that the density f_i and the tail function $\overline{F_i} = 1 - F_i$ satisfy the relation

$$f_i(x)/\overline{F_i}(x) \ge A_i/x \quad \textit{for} \quad x \ge x_0, \tag{3}$$

and there exist constants $B_i > 0$, $\alpha_i > 0$, $\beta_i > 0$ and real γ_i such that

$$\overline{F_i}(x) \ge B_i x^{\gamma_i} e^{-\alpha_i x^{\beta_i}} \quad \textit{for} \quad x \ge x_0. \tag{4}$$

If, in addition to (i) and (ii), the parameters $\beta_1, \ldots, \beta_n$ are such that $\sum_{i=1}^n \frac{1}{\beta_i} > 2$, then the product $Z_n = \xi_1 \xi_2 \cdots \xi_n$ is M-indet.

To prove Theorem 7, we need the following lemma.

Lemma 3. *Let F be a distribution on $\mathbb{R}$ such that (i) it has density f on the subset $[a, ra]$, where $a > 0$ and $r > 1$, and (ii) for some constant $A > 0$, $f(x)/\overline{F}(x) \ge A/x$ on $[a, ra]$. Then*

$$\int_a^{ra} \frac{f(x)}{x} dx \ge \left(1 - \frac{1}{r}\right) \frac{A}{1+A} \frac{\overline{F}(a)}{a}.$$

Proof. By integration by parts, we have

$$\begin{aligned} \int_a^{ra} \frac{f(x)}{x} dx &= -\int_a^{ra} \frac{d\overline{F}(x)}{x} = \frac{\overline{F}(a)}{a} - \frac{\overline{F}(ra)}{ra} - \int_a^{ra} \frac{\overline{F}(x)}{x^2} dx \\ &\ge \left(1 - \frac{1}{r}\right) \frac{\overline{F}(a)}{a} - \frac{1}{A} \int_a^{ra} \frac{f(x)}{x} dx, \end{aligned}$$

in which the last inequality is due to the monotonicity of F and the condition on the failure rate $f/\overline{F}$. Hence the required conclusion follows. $\square$

Proof of Theorem 7. We may assume, see condition (i), that f_n is the density which is decreasing in $x \ge x_0$. Then, clearly Z_n is nonnegative and its density, say h_n, can be written as follows: for $x > 0$,

$$h_n(x) = \int_0^\infty \int_0^\infty \cdots \int_0^\infty \frac{f_1(u_1)}{u_1} \frac{f_2(u_2)}{u_2} \cdots \frac{f_{n-1}(u_{n-1})}{u_{n-1}} f_n\left(\frac{x}{u_1 u_2 \cdots u_{n-1}}\right) du_1 du_2 \cdots du_{n-1}.$$

This representation shows that $h_n(x) > 0$. To prove the M-indet property of Z_n, we will show instead the finiteness of the Krein quantity of h_n. Therefore, we have to estimate the lower bound of h_n.

Since $\sum_{i=1}^{n} \frac{1}{\beta_i} > 2$, we can choose $n-1$ numbers $\theta_i \in (0,1),\ i = 1, 2, \ldots, n-1$, such that $\theta_i < \frac{1}{2\beta_i}$ and

$$1 - \frac{1}{2\beta_n} < \sum_{i=1}^{n-1} \theta_i < \min\left\{1,\ \sum_{i=1}^{n-1} \frac{1}{2\beta_i}\right\}.$$

Denote $\theta_n = 1 - \sum_{i=1}^{n-1} \theta_i$, define $\theta = \min\{\theta_1, \ldots, \theta_{n-1}, \theta_n\}$ and let $x_\theta = (2^{n-1}x_0)^{1/\theta}$. Then for each $x > x_\theta$, we take $a_i = x^{\theta_i} \geq x_0$ which implies

$$x/(2^{n-1}a_1a_2\cdots a_{n-1}) = x^{\theta_n}/2^{n-1} \geq x_0.$$

For these x and a_i, we have, by condition (i), the following:

$$\begin{aligned}
h_n(x) &\geq \int_{a_1}^{2a_1}\int_{a_2}^{2a_2}\cdots\int_{a_{n-1}}^{2a_{n-1}} \frac{f_1(u_1)}{u_1}\frac{f_2(u_2)}{u_2}\cdots\frac{f_{n-1}(u_{n-1})}{u_{n-1}} \\
&\qquad \times f_n\left(\frac{x}{u_1u_2\cdots u_{n-1}}\right) du_1du_2\cdots du_{n-1} \\
&\geq \int_{a_1}^{2a_1}\int_{a_2}^{2a_2}\cdots\int_{a_{n-1}}^{2a_{n-1}} \frac{f_1(u_1)}{u_1}\frac{f_2(u_2)}{u_2}\cdots\frac{f_{n-1}(u_{n-1})}{u_{n-1}} \\
&\qquad \times f_n\left(\frac{x}{a_1a_2\cdots a_{n-1}}\right) du_1du_2\cdots du_{n-1} \\
&= f_n\left(\frac{x}{a_1a_2\cdots a_{n-1}}\right)\prod_{i=1}^{n-1}\int_{a_i}^{2a_i}\frac{f_i(u)}{u}du.
\end{aligned}$$

Then, by Lemma 3 (with $r = 2$) and (3)–(4),

$$\begin{aligned}
h_n(x) &\geq f_n\left(\frac{x}{a_1a_2\cdots a_{n-1}}\right)\prod_{i=1}^{n-1}\frac{A_i}{2(1+A_i)}\frac{\overline{F_i}(a_i)}{a_i} \\
&\geq C\left(\frac{x}{a_1a_2\cdots a_{n-1}}\right)^{\gamma_n-1}\left(\prod_{i=1}^{n-1}a_i^{\gamma_i-1}\right) \\
&\qquad \times \exp\left[-\sum_{i=1}^{n-1}\alpha_i a_i^{\beta_i} - \alpha_n\left(\frac{x}{a_1a_2\cdots a_{n-1}}\right)^{\beta_n}\right] \\
&= Cx^{\gamma}\exp\left[-\sum_{i=1}^{n-1}\alpha_i x^{\theta_i\beta_i} - \alpha_n x^{\theta_n\beta_n}\right], \quad x > x_\theta,
\end{aligned}$$

where $C = 2^{1-n} A_n (\prod_{i=1}^{n-1} \frac{A_i}{1+A_i}) \prod_{i=1}^{n} B_i$ and $\gamma = \sum_{i=1}^{n-1} \theta_i(\gamma_i - 1) + \theta_n(\gamma_n - 1)$.

In the exponential factor above we keep separately the two terms because of their rôle when evaluating the Krein quantity on (x_θ, ∞) for h_n. Recall that this is a Stieltjes case and we have the following:

$$\mathrm{K}[h_n] = \int_{x_\theta}^{\infty} \frac{-\log h_n(x^2)}{1+x^2} dx < \infty.$$

The conclusion about the finiteness of $\mathrm{K}[h_n]$ relies essentially on the facts that $2\theta_i\beta_i < 1,\ i = 1, 2, \ldots, n-1$, and $2\theta_n\beta_n < 1$. Therefore, Z_n is M-indet by Proposition 1 in [17]. The proof is complete. □

Example 1. For illustration of how to use Theorem 7, consider the class of generalized gamma distributions. We use the notation $\xi \sim GG(\alpha, \beta, \gamma)$ if the density function of the random variable ξ is of the form

$$f(x) = cx^{\gamma-1} e^{-\alpha x^\beta}, \quad x \geq 0. \tag{5}$$

Here $\alpha, \beta, \gamma > 0$, $f(0) = 0$ if $\gamma \neq 1$, and $c = \beta\alpha^{\gamma/\beta}/\Gamma(\gamma/\beta)$ is the norming constant. We have the following statement (see also Theorem 8.4 in [18]).

Corollary 1. *Suppose $\xi_1, \ldots, \xi_n$ are n independent random variables such that $\xi_i \sim GG(\alpha_i, \beta_i, \gamma_i),\ i = 1, \ldots, n$, and let $Z_n = \xi_1 \cdots \xi_n$. Then Z_n is M-det if and only if $\sum_{i=1}^{n} \frac{1}{\beta_i} \leq 2$.*

Proof. Note that for $\xi \sim GG(\alpha, \beta, \gamma)$ defined by (5), we have two properties, namely: (a) $f(x)/\overline{F}(x) \approx \alpha\beta x^{\beta-1}$, $\overline{F}(x) \approx [c/(\alpha\beta)] x^{\gamma-\beta} e^{-\alpha x^\beta}$ as $x \to \infty$, and (b) $m_k = \alpha^{-k/\beta}\Gamma((\gamma+k)/\beta))/\Gamma(\gamma/\beta)) = \mathcal{O}(k^{k/\beta})$ as $k \to \infty$. Hence the sufficiency part follows from Theorem 1 because $\mathbf{E}[Z_n^k] = \mathcal{O}(k^{Bk})$ as $k \to \infty$, where $B = \sum_{i=1}^{n} \frac{1}{\beta_i}$, while the necessity part is a consequence of Theorem 7. □

Example 2. Consider the class of inverse Gaussian distributions. We say that $X \sim IG(\mu, \lambda)$ if the density of X is of the form

$$f(x) = \left(\frac{\lambda}{2\pi x^3}\right)^{1/2} \exp\left[-\frac{\lambda(x-\mu)^2}{2\mu^2 x}\right], \quad x > 0, \tag{6}$$

where $\mu, \lambda > 0$ and $f(0) = 0$. If $X \sim IG(\mu, \lambda)$, then it has a moment generating function. This in turn implies that the power $Y = X^2$ satisfies Hardy's condition and hence is M-det. Actually, we have that for real r, X^r is M-det if and only if $|r| \leq 2$ (see [23]). If ξ_1 and ξ_2 are two i.i.d. random variables with density (6), then the product $Z = \xi_1\xi_2$ is also M-det due to Proposition 1(iii) in [15]. The next result is for products of non-identically distributed random variables.

Corollary 2. *Let $\xi_1 \sim IG(\mu_1, \lambda_1)$, $\xi_2 \sim IG(\mu_2, \lambda_2)$ and $\eta \sim Exp(1)$ be three independent random variables. Then the following statements hold:*

(i) $Z = \xi_1 \eta$ is M-det.
(ii) $Z = \xi_1 \xi_2$ is M-det.
(iii) $Z = \xi_1 \xi_2 \eta$ is M-indet.

Proof. First, for $X \sim F = IG(\mu, \lambda)$, it can be shown (we omit the details) that the moment $\mathbf{E}[X^k] = \mathcal{O}(k^k)$ as $k \to \infty$. Second, the hazard rate function $r(x) = f(x)/\overline{F}(x) \to \lambda/(2\mu^2) > 0$ as $x \to \infty$. Third, the tail function $\overline{F}$ satisfies (4) with the exponent $\beta = 1$. With these three steps we are in a position to apply Theorems 5 and 7 to confirm the validity of (i) – (iii) as stated above. □

4. Products of Random Variables on $\mathbb{R}$

We start with two results describing relatively simple conditions on the random variables $\xi_1, \ldots, \xi_n$ in order to guarantee that their product is M-det. The results are similar to the above Theorems 5 and 6, however we remember that here we deal with the Hamburger case, so we work with the even order moments.

Theorem 8. *Suppose that the even order moments $m_{i,2k} = \mathbf{E}[\xi_i^{2k}]$, $i = 1, \ldots, n$, of the independent random variables $\xi_1, \ldots, \xi_n$ satisfy the conditions:*

$$m_{i,2k} = \mathcal{O}((2k)^{2a_i k}) \quad as \quad k \to \infty, \quad for \quad i = 1, \ldots, n,$$

where $a_1, \ldots, a_n$ are positive constants. If the parameters $a_1, \ldots, a_n$ are such that $a_1 + \cdots + a_n \leq 1$, then the product $Z_n = \xi_1 \cdots \xi_n$ is M-det.

Proof. With $m_{2k} = \mathbf{E}[Z_n^{2k}]$ we have, by the independence of ξ_i, that

$$m_{2k} = m_{1,2k} \cdots m_{n,2k} = \mathcal{O}((2k)^{2a_1 k}) \cdots \mathcal{O}((2k)^{2a_n k}) = \mathcal{O}((2k)^{2ak}) \quad \text{as} \quad k \to \infty,$$

where $a = a_1 + \cdots + a_n$. Since, by assumption, $a \leq 1$, we apply Theorem 3 to conclude the M-det property of the product Z_n. □

Theorem 9. *Suppose that the growth rates $r_1, \ldots, r_n$ of the even order moments of each of the independent random variables $\xi_1, \ldots, \xi_n$ satisfy*

$$\frac{m_{1,2(k+1)}}{m_{1,2k}} = \mathcal{O}((k+1)^{r_1}), \ \ldots, \ \frac{m_{n,2(k+1)}}{m_{n,2k}} = \mathcal{O}((k+1)^{r_n}) \quad as \quad k \to \infty,$$

where $m_{i,2k} = \mathbf{E}[\xi_i^{2k}], i = 1, \ldots, n, \ k = 1, 2, \ldots$. If the rates $r_1, \ldots, r_n$ are such that $r_1 + \cdots + r_n \leq 2$, then the product $Z_n = \xi_1 \cdots \xi_n$ is M-det.

Proof. Denoting $m_{2k} = \mathbf{E}[Z_n^{2k}]$ we have, by the independence of ξ_i, that

$$\frac{m_{2(k+1)}}{m_{2k}} = \frac{m_{1,2(k+1)}}{m_{1,2k}} \cdots \frac{m_{n,2(k+1)}}{m_{n,2k}} = \mathcal{O}((k+1)^r) \quad \text{as } k \to \infty,$$

where $r = r_1 + \cdots + r_n$. Since, by assumption, $r \leq 2$, Theorem 3 together with Remark 4 implies the M-det property of the product Z_n. □

Let us describe now conditions under which the product Z_n is M-indet.

Theorem 10. *Consider n independent random variables $\xi_i \sim F_i,\ i = 1, 2, \ldots, n$, where $n \geq 2$, and let each F_i be absolutely continuous with a symmetric density (about 0) $f_i > 0$ on $\mathbb{R}$. Assume further that the following conditions are satisfied:*
(i) At least one (or just one) of the densities $f_1(x), \ldots, f_n(x)$ is decreasing in $x \geq x_0$, where $x_0 \geq 1$ is a constant.
(ii) For each $i = 1, 2, \ldots, n$, there exists a constant $A_i > 0$ such that

$$f_i(x)/\overline{F_i}(x) \geq A_i/x \quad \textit{for} \quad x \geq x_0, \tag{7}$$

and there exist constants $B_i > 0,\ \alpha_i > 0,\ \beta_i > 0$ and real γ_i such that

$$\overline{F_i}(x) \geq B_i x^{\gamma_i} e^{-\alpha_i x^{\beta_i}} \quad \textit{for} \quad x \geq x_0. \tag{8}$$

If, in addition to the above, $\sum_{i=1}^{n} \frac{1}{\beta_i} > 1$, then the product $Z_n = \xi_1 \xi_2 \cdots \xi_n$ is M-indet.

Proof. We may assume, see condition (i), that f_n is the density which is decreasing in $x \geq x_0$. Then the density h_n of Z_n is symmetric about 0 (see, e.g., [11]) and h_n can be written as follows: for $x > 0$,

$$\begin{aligned} h_n(x) &= 2^{n-1} \int_0^\infty \int_0^\infty \cdots \int_0^\infty \frac{f_1(u_1)}{u_1} \frac{f_2(u_2)}{u_2} \cdots \frac{f_{n-1}(u_{n-1})}{u_{n-1}} \\ &\qquad \times f_n\left(\frac{x}{u_1 u_2 \cdots u_{n-1}}\right) du_1 du_2 \cdots du_{n-1}. \end{aligned}$$

Hence $h_n(x) > 0$. As in the proof of Theorem 7, we will estimate the lower bound of h_n. For completeness we give detailed arguments here.

Since $\sum_{i=1}^{n} \frac{1}{\beta_i} > 1$ by assumption, we can choose numbers $\theta_i \in (0, 1),\ i = 1, 2, \ldots, n-1$, such that $\theta_i < \frac{1}{\beta_i}$ and

$$1 - \frac{1}{\beta_n} < \sum_{i=1}^{n-1} \theta_i < \min\left\{1,\ \sum_{i=1}^{n-1} \frac{1}{\beta_i}\right\}.$$

Denote $\theta_n = 1 - \sum_{i=1}^{n-1} \theta_i$, define $\theta = \min\{\theta_1, \theta_2, \ldots, \theta_{n-1}, \theta_n\}$ and let $x_\theta =$

$(2^{n-1}x_0)^{1/\theta}$. By taking $x > x_\theta$ and $a_i = x^{\theta_i}$, we obtain the following:

$$\begin{aligned} h_n(x) &\geq 2^{n-1} \int_{a_1}^{2a_1} \int_{a_2}^{2a_2} \cdots \int_{a_{n-1}}^{2a_{n-1}} \frac{f_1(u_1)}{u_1} \frac{f_2(u_2)}{u_2} \cdots \frac{f_{n-1}(u_{n-1})}{u_{n-1}} \\ &\qquad \times f_n\left(\frac{x}{u_1 u_2 \cdots u_{n-1}}\right) du_1 du_2 \cdots du_{n-1} \\ &\geq 2^{n-1} \int_{a_1}^{2a_1} \int_{a_2}^{2a_2} \cdots \int_{a_{n-1}}^{2a_{n-1}} \frac{f_1(u_1)}{u_1} \frac{f_2(u_2)}{u_2} \cdots \frac{f_{n-1}(u_{n-1})}{u_{n-1}} \\ &\qquad \times f_n\left(\frac{x}{a_1 a_2 \cdots a_{n-1}}\right) du_1 du_2 \cdots du_{n-1} \\ &= 2^{n-1} f_n\left(\frac{x}{a_1 a_2 \cdots a_{n-1}}\right) \prod_{i=1}^{n-1} \int_{a_i}^{2a_i} \frac{f_i(u)}{u} du. \end{aligned}$$

Applying Lemma 3 and the above conditions (7) and (8), we further have

$$\begin{aligned} h_n(x) &\geq 2^{n-1} f_n\left(\frac{x}{a_1 a_2 \cdots a_{n-1}}\right) \prod_{i=1}^{n-1} \frac{A_i}{2(1+A_i)} \frac{\overline{F_i}(a_i)}{a_i} \\ &\geq C \left(\frac{x}{a_1 a_2 \cdots a_{n-1}}\right)^{\gamma_n - 1} \left(\prod_{i=1}^{n-1} a_i^{\gamma_i - 1}\right) \\ &\qquad \times \exp\left[-\sum_{i=1}^{n-1} \alpha_i a_i^{\beta_i} - \alpha_n \left(\frac{x}{a_1 a_2 \cdots a_{n-1}}\right)^{\beta_n}\right] \\ &= C x^{\tilde{\gamma}} \exp\left[-\sum_{i=1}^{n-1} \alpha_i x^{\theta_i \beta_i} - \alpha_n x^{\theta_n \beta_n}\right], \quad x > x_\theta. \end{aligned}$$

Here $C = A_n(\prod_{i=1}^{n-1} \frac{A_i}{1+A_i}) \prod_{i=1}^{n} B_i$ and $\tilde{\gamma} = \sum_{i=1}^{n-1} \theta_i(\gamma_i - 1) + \theta_n(\gamma_n - 1)$.

The next step is to evaluate the Krein quantity on $|x| > x_\theta$ for h_n in this Hamburger case:

$$\mathrm{K}[h_n] = 2 \int_{x_\theta}^{\infty} \frac{-\log h_n(x)}{1+x^2} dx < \infty$$

because $\theta_i \beta_i < 1$ for each $i = 1, 2, \ldots, n-1$ and $\theta_n \beta_n < 1$. Hence, the product Z_n is M-indet by Theorem 2.2 in [19] (see also [17]). □

Example 3. We now apply Theorem 10 to the product of double generalized gamma random variables. We write $\xi \sim DGG(\alpha, \beta, \gamma)$ if ξ is a random variable in $\mathbb{R}$ with density function of the form

$$f(x) = c|x|^{\gamma - 1} e^{-\alpha |x|^\beta}, \quad x \in \mathbb{R}. \tag{9}$$

Here $\alpha, \beta, \gamma > 0$, $f(0) = 0$ if $\gamma \neq 1$, and $c = \beta\alpha^{\gamma/\beta}/(2\Gamma(\gamma/\beta))$ is a norming constant.

Corollary 3. *Suppose that $\xi_1, \ldots, \xi_n$ are n independent random variables, and let $\xi_i \sim DGG(\alpha_i, \beta_i, \gamma_i)$, $i = 1, 2, \ldots, n$. Then the product $Z_n = \xi_1 \cdots \xi_n$ is M-det if and only if $\sum_{i=1}^n \frac{1}{\beta_i} \leq 1$.*

Proof. Note that for the moment $m_{2k} = \mathbf{E}[\xi^{2k}]$ of $\xi \sim DGG(\alpha, \beta, \gamma)$, see (9), we have the following relation: $m_{2k} = \mathcal{O}((2k)^{2k/\beta})$ as $k \to \infty$. Thus, the sufficiency part is exactly Theorem 10 in [25]. The same statement can also be proved by Theorem 8 above. Finally, the necessity part follows from Theorem 10 (we may redefine $f(0)$ to be a positive number if necessary). □

5. The Mixed Case

For completeness of our study we need to consider products of both types of random variables, nonnegative ones and real ones. Since such a 'mixed' product takes values in $\mathbb{R}$, this is a Hamburger case, so we can formulate results similar to Theorems 8 and 9. Since the conditions, the statements and the arguments are almost as in these two theorems, we do not give details. Instead, we suggest now a result in which the 'mixed' product $Z_n = \xi_1 \cdots \xi_n$ is M-indet.

Theorem 11. *Given are n independent random variables, such that the 'first' group, $\xi_1, \ldots, \xi_{n_0}$, consists of nonnegative variables, while the variables in the 'second' group, $\xi_{n_0+1}, \ldots, \xi_n$, are all with values in $\mathbb{R}$, where $1 \leq n_0 < n$. We assume that all $\xi_i \sim F_i$, $i = 1, \ldots, n$, are absolutely continuous; denote their densities by f_i, $i = 1, \ldots, n$. Assume further that $f_i(x) > 0$ on $(0, \infty)$, $i = 1, \ldots, n_0$, while $f_j(x) > 0$ on $\mathbb{R}$, $j = n_0 + 1, \ldots, n$, and are symmetric. We require also the following conditions:*
(i) At least one of the densities $f_j(x)$, $j = n_0 + 1, \ldots, n$, is decreasing in $x \geq x_0$, where $x_0 \geq 1$ is a constant.
(ii) For each $i = 1, 2, \ldots, n$, there exist a constant $A_i > 0$ such that

$$f_i(x)/\overline{F_i}(x) \geq A_i/x \quad \text{for} \quad x \geq x_0,$$

and there exist constants $B_i > 0$, $\alpha_i > 0$, $\beta_i > 0$ and real γ_i such that

$$\overline{F_i}(x) \geq B_i x^{\gamma_i} e^{-\alpha_i x^{\beta_i}} \quad \text{for} \quad x \geq x_0.$$

If, in addition to (i) and (ii), the parameters β_i are such that $\sum_{i=1}^n \frac{1}{\beta_i} > 1$, then the product $Z_n = \xi_1 \cdots \xi_n$ is M-indet.

Proof. The proof is similar to that of Theorem 10 and can be omitted; the only change is to replace the coefficient 2^{n-1} in the integral form of h_n by 2^{n-n_0-1}. □

As an application of Theorem 11 we derive below two interesting corollaries.

Corollary 4. *Consider two independent random variables, ξ and η, where $\xi \sim Exp(1)$ and $\eta \sim \mathcal{N}(0,1)$ (standard normal). Then $Z = \xi\,\eta$ is M-indet.*

In a similar way we arrive also at the following statement.

Corollary 5. *(i) The product of two independent random variables, one chi-square and one normal, is M-indet.*
(ii) The product of two independent random variables, one inverse Gaussian and one normal, is M-indet.

Acknowledgements

The work of GDL was partly supported by the Ministry of Science and Technology of Taiwan (ROC) under Grant NSC 102-2118-M-001-008-MY2.

JS was supported by the International Laboratory of Quantitative Finance (Higher School of Economics, Moscow) under Grant No. 14.12.31.0007 to visit Moscow, March 2014, when a part of this work was completed. The work was finalized during the visit of JS to Academia Sinica (Taipei, Taiwan, ROC), May 2014, with a support from the Emeritus Fellowship provided by Leverhulme Trust (UK).

References

[1] Berg, C.: On powers of Stieltjes moment sequences, I. J. Theoret. Probab. **18**, 871–889 (2005)

[2] Bisgaard, T.M., Sasvári, Z.: Characteristic Functions and Moment Sequences: Positive Definiteness in Probability. Nova Sci. Publishers, Huntington (NY) (2000)

[3] Bondesson, L.: A class of probability distributions that is closed with respect to addition as well as multiplication of independent random variables. J. Theoret. Probab. doi:10.1007/s10959-013-0523-y (2013)

[4] Carmona, R., Molchanov, S.A.: Intermittency and phase transitions for some particle systems in random media. In: Elworthy, K.D., Ikeda, N. (eds.) Asymptotic Problems in Probability Theory: Stochastic Models and Diffusions on Fractals (Sanda/Kyoto, 1990), pp. 15–36, Pitman Research Notes in Mathematics Series **283**. Longman Science & Technology, Harlow (1993)

[5] Chen, Y., Karagiannidis, G.K., Lu, H., Cao, N.: Novel approximations to the statistics of products of independent random variables and their applications in wireless communications. IEEE Trans. on Vehicular Tech. **61**, 443–454 (2012)

[6] Cline, D.B.H., Samorodnitsky, G.: Subexponentiality of the product of independent random variables. Stochastic Process. Appl. **49**, 75–98 (1994)

[7] Davis, R.A., Mikosch, T.: Probabilistic properties of stochastic volatility models. In: Andersen, T.G., Davis, R.A., Kreiss, J.-P., Mikosch, T. (eds.) Handbook of Financial Time Series, pp. 255–267. Springer, New York (2009)

[8] de Abreu, G.T.F.: On the moment determinance and random mixture of Nakagami-m variates. IEEE Trans. on Communications **58**, 2561–2575 (2010)

[9] Frisch, U., Sornette, D.: Extreme deviations and applications. J. Physique Serie I **7**, 1155–1171 (1997)

[10] Galambos, J., Simonelli, I.: Products of Random Variables: Applications to Problems of Physics and to Arithmetical Functions. Marcel Dekker, New York (2004)

[11] Hamedani, G.G., Walter, G.G.: On the product of symmetric random variables. Statist. Probab. Lett. **3**, 251–253 (1985) Erratum: ibid **4** (1986), 51.

[12] Hashorva, E., Pakes, A.G., Tang, Q.: Asymptotics of random contractions. Insurance Math. Econom. **47**, 405–414 (2010)

[13] Lévy, P.: Esquisse d'une théorie de la multiplication des variables aléatoires. Annales Sci. de l'É.N.S. 3e Ser. **76**, 59–82 (1959)

[14] Lin, G.D.: On the moment problems. Statist. Probab. Lett. **35**, 85–90 (1997) Erratum: ibid **50** (2000), 205.

[15] Lin, G.D., Stoyanov, J.: Moment determinacy of powers and products of nonnegative random variables. J. Theoret. Probab. In press. doi:10.1007/s10959-014-0546-z (2014)

[16] Lomnicki, Z.A.: On the distribution of products of random variables. J. Roy. Statist. Soc. Ser. B **29**, 513–524 (1967)

[17] Pakes, A.G.: Remarks on converse Carleman and Krein criteria for the classical moment problem. J. Austral. Math. Soc. **71**, 81–104 (2001)

[18] Pakes, A.G.: On generalized stable and related laws. J. Math. Anal. Appl. **411**, 201–222 (2014)

[19] Pedersen, H.L.: On Krein's theorem for indeterminacy of the classical moment problem. J. Approx. Theory **95**, 90–100 (1998)

[20] Penson, K.A., Blasiak, P., Duchamp, G.H.E., Horzela, A., Solomon, A.I.: On certain non-unique solutions of the Stieltjes moment problem. Discr. Math. Theoret. Comput. Sci. **12**, 295–306 (2010)

[21] Shimura, T.: The product of independent random variables with regularly varying tails. Acta Appl. Math. **63**, 411–432 (2000)

[22] Springer, M.D.: The Algebra of Random Variables. Wiley, New York (1979)

[23] Stoyanov, J.: Inverse Gaussian distribution and the moment problem. J. Appl. Statist. Sci. **9**, 61–71 (1999)

[24] Stoyanov, J., Lin, G.D.: Hardy's condition in the moment problem for probability distributions. Theory Probab. Appl. **57**, 811–820 (2012) (SIAM edition: **57**, 699–708 (2013))

[25] Stoyanov, J., Lin, G.D., DasGupta, A.: Hamburger moment problem for powers and products of random variables. J. Statist. Plann. Inference **154**, 166–177 (2014)

[26] Tang, Q.: From light tails to heavy tails through multiplier. Extremes **11**, 379–391 (2008)

[27] Whittaker, E.T., Watson, G.N.: A Course of Modern Analysis. 4th edn. Cambridge University Press, Cambridge (1927)